arXiv:1404.6884v1 [math.NT] 28 Apr 2014

POLYNOMIAL MAPS ON VECTOR SPACES OVER A FINITE FIELD

MICHIEL KOSTERS

Abstract. Let l be a finite field of cardinality q and let n be in $\mathbf{Z}_{\geq 1}$. Let $f_1, \ldots, f_n \in l[x_1, \ldots, x_n]$ not all constant and consider the evaluation map $f = (f_1, \ldots, f_n) \colon l^n \to l^n$. Set $\deg(f) = \max_i \deg(f_i)$. Assume that $l^n \setminus f(l^n)$ is not empty. We will prove

$$|l^n \setminus f(l^n)| \geq \frac{n(q-1)}{\deg(f)}.$$

This improves previous known bounds.

1. Introduction

The main result of [MWW12] is the following theorem.

Theorem 1.1. *Let l be a finite field of cardinality q and let n be in $\mathbf{Z}_{\geq 1}$. Let $f_1, \ldots, f_n \in l[x_1, \ldots, x_n]$ not all constant and consider the map $f = (f_1, \ldots, f_n) \colon l^n \to l^n$. Set $\deg(f) = \max_i \deg(f_i)$. Assume that $l^n \setminus f(l^n)$ is not empty. Then we have*

$$|l^n \setminus f(l^n)| \geq \min\left\{\frac{n(q-1)}{\deg(f)}, q\right\}.$$

We refer to [MWW12] for a nice introduction to this problem including references and historical remarks. The proof in [MWW12] relies on p-adic liftings of such polynomial maps. We give a proof of a stronger statement using different techniques.

Theorem 1.2. *Under the assumptions of Theorem 1.1 we have*

$$|l^n \setminus f(l^n)| \geq \frac{n(q-1)}{\deg(f)}.$$

We deduce the result from the case $n = 1$ by putting a field structure k on l^n and relate the k-degree and the l-degree. We prove the result $n = 1$ in a similar way as in [Tur95].

2. Degrees

Let l be a finite field of cardinality q and let V be a finite dimensional l-vector space. By $V^\vee = \mathrm{Hom}(V, l)$ we denote the dual of V. Let $v_1, \ldots, v_f$ be a basis of V. By $x_1, \ldots, x_f$ we denote its dual basis in $V^\vee$, that is, x_i is the map which sends v_j to δ_{ij}. Denote by $\mathrm{Sym}_l(V^\vee)$ the symmetric algebra of $V^\vee$ over l. We have an isomorphism $l[x_1, \ldots, x_f] \to \mathrm{Sym}_l(V^\vee)$ mapping x_i to x_i. Note that $\mathrm{Map}(V, l) = l^V$

Date: April 29, 2014.

2010 *Mathematics Subject Classification.* 11T06.

This article is part of my PhD thesis written under the supervision of Hendrik Lenstra.

is a commutative ring under the coordinate wise addition and multiplication and it is a l-algebra. The linear map $V^\vee \to \mathrm{Map}(V,l)$ induces by the universal property of $\mathrm{Sym}_l(V^\vee)$ a ring morphism $\varphi\colon \mathrm{Sym}_l(V^\vee) \to \mathrm{Map}(V,l)$. When choosing a basis, we have the following commutative diagram, where the second horizontal map is the evaluation map and the vertical maps are the natural isomorphisms:

$$\begin{array}{ccc} \mathrm{Sym}(V^\vee) & \xrightarrow{\varphi} & \mathrm{Map}(V,l) \\ \uparrow & & \uparrow \\ l[x_1,\ldots,x_n] & \longrightarrow & \mathrm{Map}(l^n,l). \end{array}$$

Lemma 2.1. *The map φ is surjective. After a choice of a basis as above the kernel is equal to $(x_i^q - x_i : i = 1,\ldots,f)$ and every $f \in \mathrm{Map}(V,l)$ has a unique representative $\sum_{m=(m_1,\ldots,m_n)\colon\ 0\leq m_i\leq q-1} c_m x_1^{m_1}\cdots x_n^{m_n}$ with $c_m \in l$.*

Proof. After choosing a basis, we just consider the map

$$l[x_1,\ldots,x_f] \to \mathrm{Map}(l^n,l).$$

For $c = (c_1,\ldots,c_f) \in l^f$ set

$$f_c = \prod_i (1-(x_i-c_i)^q).$$

For $c' \in l^f$ we have $f_c(c') = \delta_{cc'}$. With these building blocks one easily shows that φ is surjective.

For $i \in \{1,2,\ldots,f\}$ the element $x_i^q - x_i$ is in the kernel of φ. This shows that modulo the kernel any $g \in l[x_1,\ldots,x_f]$ has a representative

$$f = \sum_{m=(m_1,\ldots,m_r)\colon\ 0\leq m_i\leq q-1} c_m x_1^{m_1}\cdots x_r^{m_n}.$$

The set of such elements has cardinality q^{q^r}. As $\#\,\mathrm{Map}(V,l) = q^{q^r}$, we see that the kernel is $(x_i^q - x_i : i = 1,\ldots,r)$. Furthermore, any element has a unique representative as described above. □

Note that $\mathrm{Sym}_l(V^\vee)$ is a graded l-algebra where we say that 0 has degree $-\infty$. For $f \in \mathrm{Map}(V,l)$ we set

$$\deg_l(f) = \min\left(\deg(g) : \varphi(g) = f\right).$$

Note that $\deg_l(f_1+f_2) \leq \max(\deg_l(f_1),\deg_l(f_2))$, with equality if the degrees are different. In practice, if $f \in l[x_1,\ldots,x_n]$, then $\deg_l(f)$ is calculated as follows: for all i replace x_i^q by x_i until $\deg_{x_i}(f) < q$. Then the degree is the total degree of the remaining polynomial.

Let W be a finite dimensional l-vector space. Then we have $\mathrm{Map}(V,W) = W \otimes_l \mathrm{Map}(V,l)$. For $f \in \mathrm{Map}(V,W)$ we set

$$\deg_l(f) = \max\left(\deg_l(g\circ f) : g \in W^\vee\right).$$

If $g_1,\ldots,g_n$ is a basis of $W^\vee$, then $\deg_l(f) = \max\left(\deg_l(g_i\circ f) : i = 1,\ldots,n\right)$. This follows from the identity $\deg_l(\sum_i c_i g_i) \leq \max(\deg_l(g_i))$ for $c_i \in l$. Note that the degree is bounded above by $(q-1)\cdot\dim_l(V)$.

For $i \in \mathbf{Z}_{\geq 0}$ and a subset S of $\mathrm{Sym}_l(V^\vee)$ we set

$$S_l^i = \mathrm{Span}_l(s_1\cdot\ldots\cdot s_i : s_i \in S) \in \mathrm{Sym}_l(V^\vee).$$

Lemma 2.2. *Let $f \in \mathrm{Map}(V,W)$. For $i \in \mathbf{Z}_{\geq 0}$ one has:* $\deg_f \leq i \iff f \in W \otimes_l (l + V^\vee)^i_l$.

Proof. Suppose first that $W = l$. The proof comes down to showing the following identity for $i \in \mathbf{Z}_{\geq 0}$:

$$l + V^\vee + \ldots + (V^\vee)^i_l = (l + V^\vee)^i_l.$$

The general case follows easily. □

3. Relations between degrees

Let k be a finite field and let l be a subfield of cardinality q. Set $h = [k : l]$. Let V and W be finite dimension k-vector spaces. Let $f \in \mathrm{Map}(V,W)$. In this section we will describe the relation between the k-degree and the l-degree.

Let us first assume that $W = k$. Let $v_1, \ldots, v_r$ be a basis of V over k. Let $R = k[x_1, \ldots, x_r]/(x_1^{q^h} - x_1, \ldots, x_r^{q^h} - x_r)$. We have the following diagram where all morphisms are ring morphisms. Here ψ is the map discussed before, τ is the natural isomorphism, $\overline{\varphi}$ is the isomorphism discussed before, and σ is the isomorphism, depending on the basis, discussed above.

$$\begin{array}{ccc} k \otimes_l \mathrm{Map}(V,l) & \xrightarrow{\quad\tau\quad} & \mathrm{Map}(V,k) \\ \uparrow{\scriptstyle \psi} & & \uparrow{\scriptstyle \overline{\varphi}} \\ k \otimes_l \mathrm{Sym}_l(\mathrm{Hom}_l(V,l)) & & \mathrm{Sym}_k(\mathrm{Hom}_k(V,k))/\ker(\varphi) \xrightarrow{\sigma} R. \end{array}$$

Consider the ring morphism $\rho = \sigma \circ \overline{\varphi}^{-1} \circ \tau \circ \psi \colon k \otimes_l \mathrm{Sym}_l(\mathrm{Hom}_l(V,l)) \to R$. Lemma 2.2 suggest that to compare degrees, we need to find

$$\rho(k \otimes_l (l + \mathrm{Hom}_l(V,l))^i_l).$$

The following lemma says that it is enough to find $k + k \otimes_l \mathrm{Hom}_l(V,l)$.

Lemma 3.1. *For $i \in \mathbf{Z}_{\geq 0}$ we have the following equality in $k \otimes_l \mathrm{Sym}_l(V)$:*

$$k \otimes_l (l + \mathrm{Hom}_l(V,l))^i_l = (k + k \otimes_l \mathrm{Hom}_l(V,l))^i_k .$$

Proof. Both are k-vector spaces and the inclusions are not hard to see. □

The following lemma identifies $k + k \otimes_l \mathrm{Hom}_l(V,l)$.

Lemma 3.2. *One has*

$$\rho(k + k \otimes_l \mathrm{Hom}_l(V,l)) = \mathrm{Span}_k \left(\{ x_j^{q^s} : 1 \leq j \leq r,\ 0 \leq s < h \} \sqcup \{1\} \right).$$

Proof. Note that $\tau \circ \phi(k + k \otimes_l \mathrm{Hom}_l(V,l)) = k \oplus \mathrm{Hom}_l(V,k) \subseteq \mathrm{Map}(V,k)$. Note that

$$\sigma^{-1} \left(\mathrm{Span}_k \left(\{ x_j^{q^s} : 1 \leq j \leq r,\ 0 \leq s < h \} \right) \right) \subseteq \mathrm{Hom}_l(V,k).$$

As both sets have dimension $\dim_l(V) = r \cdot h$ over k, the result follows. □

For $m, n \in \mathbf{Z}_{\geq 1}$ we set $s_m(n)$ to be the sum of the digits of n in base m.

Lemma 3.3. *Let $m \in \mathbf{Z}_{\geq 2}$ and $n, n' \in \mathbf{Z}_{\geq 0}$. Then the following hold:*

i. $s_m(n + n') \leq s_m(n) + s_m(n')$;
ii. *Suppose $n = \sum_i c_i m^i$, $c_i \geq 0$. Then we have $\sum_i c_i \geq s_m(n)$ with equality iff for all i we have $c_i < m$.*

Proof. i. This is well-known and left to the reader.

ii. We give a proof by induction on n. For $n = 0$ the result is correct. Suppose first that $n = c_s m^s$ and assume that $c_s \geq m$. Then we have $n = (c_s - m)m^s + m^{s+1}$. By induction and i we have

$$c_s > c_s - m + 1 \geq s_m((c_s - m)m^s) + s_m(m^{s+1}) \geq s_m(c_s m^s).$$

In general, using i, we find

$$\sum_i c_i \geq \sum_i s_m(c_i m^i) \geq s_m(n).$$

Also, one easily sees that one has equality iff all c_i are smaller than m. □

Proposition 3.4. *Let $f \in k[x_1, \ldots, x_r]$ nonzero with the degree in all x_i of all the monomials less than q^h. Write $f = \sum_{s=(s_1,\ldots,s_r)} c_s x_1^{s_1} \cdots x_r^{s_r}$. Then the l-degree of $\tau^{-1} \circ \overline{\varphi}(\overline{f}) \in k \otimes_l \mathrm{Map}(V, l)$ is equal to*

$$\max\{s_q(s_1) + \ldots + s_q(s_r) : s = (s_1, \ldots, s_r) \text{ s.t. } c_s \neq 0\}.$$

Proof. Put $g = \tau^{-1} \circ \overline{\varphi} \circ \sigma^{-1}(\overline{f})$. From Lemma 2.2, Lemma 3.1 and Lemma 3.2 we obtain the following. Let $i \in \mathbf{Z}_{\geq 0}$. Then $\deg_l(g) \leq i$ iff

$$\begin{aligned} g \in \rho(k \otimes_l (l + \mathrm{Hom}_l(V, l))_l^i) &= \rho((k + k \otimes_l \mathrm{Hom}_l(V, l))_k^i) \\ &= \left(\mathrm{Span}_k\left(\{x_j^{q^s} : 1 \leq j \leq r,\ 0 \leq s < h\} \sqcup \{1\}\right)\right)_k^i. \end{aligned}$$

The result follows from Lemma 3.3. □

The case for a general W just follows by decomposing W into a direct sum of copies of k and then taking the maximum of the corresponding degrees.

4. Proof of main theorem

Lemma 4.1. *Let $m, q, h \in \mathbf{Z}_{>0}$ and suppose that $q^h - 1 | m$. Then we have: $s_q(m) \geq h(q-1)$.*

Proof. We do a proof by induction on m.

Suppose that $m < q^h$. Then $m = q^h - 1$ and we have $s_q(m) = h(q-1)$.

Suppose $m \geq q^h$. Write $m = m_0 q^h + m_1$ with $0 \leq m_1 < q^h$ and $m_0 \geq 1$. We claim that $q^h - 1 | m_0 + m_1$. Note that $m_0 + m_1 \equiv m_0 q^h + m_1 \equiv 0 \pmod{q^h - 1}$. Then by induction we find

$$s_q(m) = s_q(m_0) + s_q(m_1) \geq s_q(m_0 + m_1) \geq h(q-1).$$

□

Lemma 4.2. *Let k be a finite field of cardinality q'. Let $R = k[X_a : a \in k]$ and consider the action of k^* on R given by*

$$\begin{aligned} k^* &\mapsto \mathrm{Aut}_{k-\mathrm{alg}}(R) \\ c &\mapsto (X_a \mapsto X_{ca}). \end{aligned}$$

Let $F \in R$ fixed by the action of k^* with $F(0, \ldots, 0) = 0$ and such that the degree of no monomial of F is a multiple of $q' - 1$. Then for $w = (a)_a \in k^k$ we have $F(w) = 0$.

Proof. We may assume that F is homogeneous with $d = \deg(F)$ which is not a multiple of $q' - 1$. Take $\lambda \in k^*$ a generator of the cyclic group. As F is fixed by k^* we find:

$$F(w) = F(\lambda w) = \lambda^d F(w).$$

As $\lambda^d \neq 1$, we have $F(w) = 0$ and the result follows. □

Finally we can state and prove a stronger version of Theorem 1.2.

Theorem 4.3. *Let k be a finite field. Let $l \subseteq k$ be a subfield with $[k : l] = h$ and let V be a finite dimensional k-vector space. Let $f \in \mathrm{Map}(V, V)$ be a non-constant and non-surjective map. Then f misses at least*

$$\frac{\dim_k(V) \cdot h \cdot (\#l - 1)}{\deg_l(f)}$$

values.

Proof. Set $\#l = q$. Put a k-linear multiplication on V such that it becomes a field. This allows us reduce to the case where $V = k$. Assume $V = k$. After shifting we may assume $f(0) = 0$. Put an ordering $\leq$ on k. In $k[T]$ we have

$$\prod_{a \in k}(1 - f(a)T) = 1 - \sum_a f(a)T + \sum_{a<b} f(a)f(b)T^2 - \ldots = \sum_i a_i T^i.$$

For $1 \leq i < \frac{h(q-1)}{\deg_l(f)}$ we claim that $a_i = 0$. Put $f_0 \in k[x]$ a polynomial of degree at most $q^h - 1$ inducing $f \colon k \to k$. Consider $g_i = \prod_{a_1 < \ldots < a_i} f_0(X_{a_1}) \cdots f_0(X_{a_i})$ in $k[X_a : a \in k]$, which is fixed by k^*. We have a map

$$\varphi \colon k[X_a : a \in k] \to \mathrm{Map}(k^k, k).$$

Proposition 3.4 gives us that $\deg_l(\varphi(g_i)) = i \deg_l(f) < h(q-1)$. We claim that there is no monomial in g_i with degree a multiple of $q^h - 1$. Indeed, suppose that there is a monomial $cX_{a_1}^{r_1} \cdots X_{a_i}^{r_i}$ $(c \neq 0)$ in g_i (note that not all r_i are zero) and suppose that $q^h - 1 | \sum_i r_i$. Then by Lemma 4.1 and Proposition 3.4 we have

$$h(q-1) \leq s_q(\sum_j r_j) \leq \sum_j s_q(r_j) \leq i \cdot \deg_l(f) < h(q-1),$$

contradiction. Hence we can apply Lemma 4.2 to conclude that $a_i = 0$.

Hence we conclude

$$\prod_{a \in k}(1 - f(a)T) \equiv 1 \pmod{T^{\lceil \frac{h(q-1)}{\deg_l(f)} \rceil}}.$$

Similarly, for the identity function of l-degree 1, we conclude

$$\prod_{a \in k}(1 - aT) \equiv 1 \pmod{T^{\lceil \frac{h(q-1)}{\deg_l(f)} \rceil}}.$$

Combining this gives:

$$\begin{aligned}\prod_{a\in k\setminus f(k)}(1-aT) &= \frac{\prod_{a\in k}(1-aT)}{\prod_{b\in f(k)}(1-bT)}\\ &\equiv \frac{\prod_{a\in k}(1-aT)}{\prod_{b\in f(k)}(1-bT)}\cdot\prod_{c\in k}(1-f(c)T) \pmod{T^{\lceil\frac{h(q-1)}{\deg_l(f)}\rceil}}\\ &\equiv \prod_{b\in f(k)}(1-bT)^{\#f^{-1}(b)-1} \pmod{T^{\lceil\frac{h(q-1)}{\deg_l(f)}\rceil}}.\end{aligned}$$

Note that the polynomials $\prod_{a\in k\setminus f(k)}(1-aT)$ and $\prod_{b\in f(k)}(1-bT)^{\#f^{-1}(b)-1}$ have degree bounded by $s=k\setminus f(k)$ and are different since $s\geq 1$. But this implies that $s\geq\frac{h(q-1)}{\deg_l(f)}$. □

Remark 4.4. Different l in Theorem 4.3 may give different lower bounds.

5. Examples

In this section we will give examples which meet the bound from Theorem 1.2.

Example 5.1 ($n=\deg(f)$)**.** Let l be a finite field of cardinality q. In this example we will show that for $n,d\in\mathbf{Z}_{\geq 2}$ there are functions $f_1,\ldots,f_n\in l[x_1,\ldots,x_n]$ such that the maximum of the degrees is equal to d such that the induced map $f\colon l^n\to l^n$ satisfies $|l^n\setminus f(l^n)|=\frac{n(q-1)}{d}=q-1$. For $i=1,\ldots,n-1$ set $f_i=x_i$. Let l_{n-1} be the unique extension of l of degree $n-1$. Let $v_1,\ldots,v_{n-1}$ be a basis of l_{n-1} over l. Then $g=\mathrm{Norm}_{l_{n-1}/l}(x_1v_1+\ldots+x_{n-1}v_{n-1})$ is a homogeneous polynomial of degree $n-1$ in $x_1,\ldots,x_{n-1}$. Put $f_n=x_n\cdot g$. As the norm of a nonzero element is nonzero, one easily sees that $l^n\setminus f(l^n)=\{0\}\times\ldots\{0\}\times l^*$ has cardinality $q-1$.

Example 5.2 ($n=\frac{\deg(f)}{q-1}$)**.** Let l be a finite field and let $n\in\mathbf{Z}_{\geq 1}$. Let $f_1,\ldots,f_n\in l[x_1,\ldots,x_n]$ such that the combined map $f\colon l^n\to l^n$ satisfies $|l^n\setminus f(l^n)|=1$ (Lemma 2.1). From Theorem 1.2 and the upper bound $n(q-1)$ for the degree we deduce that $\deg(f)=n(q-1)$.

References

[MWW12] Gary L. Mullen, Daqing Wan, and Qiang Wang. Value sets of polynomial maps over finite fields. `http://arxiv.org/abs/1210.8119`, 2012. preprint.

[Tur95] Gerhard Turnwald. A new criterion for permutation polynomials. *Finite Fields Appl.*, 1(1):64–82, 1995.

Mathematisch Instituut P.O. Box 9512 2300 RA Leiden the Netherlands
E-mail address: `mkosters@math.leidenuniv.nl`
URL: `www.math.leidenuniv.nl/~mkosters`